

FDCSUSYDecay: A MSSM Decay Package

Wei Qi, J.X. Wang

*Institute of High Energy Physics, the Chinese Academy of Sciences,
P.O.Box 918(4), Beijing 100049, China*

February 2, 2008

Abstract

FDCSUSYDecay is a FORTRAN program package generated by FDC (Feynman Diagram Calculation) system fully automatically. It is dedicated to calculate at tree-level all the possible 2-body decays of SUSY and Higgs particles in the Minimal Supersymmetric extension of the Standard Model (MSSM). The format of its output files complies with SUSY Les Houches Accord and can be easily imported by other packages.

PACS: 02.70-c; 12.60 Jv

Program Summary

Program Title: FDCSUSYDecay (Version 1.00)

Journal Reference:

Catalogue identifier:

Keywords: SUSY decay, MSSM, FDC

PACS: 02.70-c; 12.60 Jv

Operating system: Linux

Programming language: FORTRAN 77

External libraries: CERNLIB 2003 (or up)

Distribution format: tar gzip file

Size of the compressed distribution file: 622,560 Bytes.

Classification: 11.1

Nature of physical problem: This package can calculate all the possible SUSY particle and Higgs 2-body decay width and branch ratio at tree-level in the MSSM model.

Method of solution: By running FDC, the Feynman rules for the MSSM model are generated, all the decay widths are calculated analytically and corresponding FORTRAN codes are generated for this package.

1 Generality

With the progress in high energy physics, it is obvious that an automatic calculation of the physical processes by the perturbative quantum field theory becomes

very useful and important. There are many projects for this purpose such as CompHEP[1], Grace[2] and so on. FDC (Feynman Diagram Calculation) [3] is one of such kind projects. It is mainly written in REDUCE and RLisp. As designed to be, it can conveniently implement physical models from the first principle, then construct their *Lagrangians*, deduce the Feynman rules and generate FORTRAN codes for calculations of physical quantities such as decay width, cross-section and matrix-element event generator. The MSSM model is implemented in FDC[4] and the generated results could be found on FDC homepage¹. To check the results such as the mixing matrices, Feynman rules, as well as to provide an application of FDC, we developed a program to automatically generate the FORTRAN packages `FDCSUSYDecay` calculating SUSY particle decays and `FDCEvent@LHC` [5] which is a matrix-element event generator of the MSSM related processes at LHC.

There are many program packages which can calculate the SUSY particle decays, such as multi-purpose packages PYTHIA, ISAJET and SUSYGEN or dedicated packages HDecay (which is special for Higgs decay in the SM and MSSM) [6] and SDecay [7], but the codes of these packages are manually typed. However, `FDCSUSYDecay` is a FORTRAN program package which is fully automatically generated by FDC. In this way many uncared errors can be systematically debugged because these artificial errors in the model would affect a series of the resulting Feynman vertices and thus the corresponding decay equations which could be found easily. All the possible 2-body decay processes (in total 543 processes) for SUSY and Higgs particles in the MSSM model have been included. This package can calculate the decay width and branch ratio at Snowmass benchmark points 1-9 [8] (Tab. 1) or any user-specified parameter configuration. The format of its output files complies with the SUSY Les Houches Accord (SLHA)[9] and thus can be imported by other programs conveniently.

Although presently only 2-body SUSY decays are implemented in this package, the FORTRAN codes for any 3-body decay can be generated by FDC. Since the important decay channels can not be distinguished from neglectable ones before the MSSM parameters are chosen and to include all 3-body decays will make the length of the generated FORTRAN codes out of control, the 3-body decays are not included in the package. Alternatively, The package with needed 3-body channels can be generated by FDC when the MSSM parameters fixed.

The package can be used in two modes. The first mode is called the high-energy scale mode. In this mode, users need to input the parameters in high-energy scale scenarios (mSUGRA, GMSB and AMSB) or choose from nine Snowmass benchmark points (which have been fed into this package hard-wiredly). Then the program obtains all the parameters at SUSY-breaking scale through interface with ISAJET [10] and then calculates everything such as masses, mixing matrices, coupling constants and decay width. The second mode is called low-energy scale mode, in which users need to input manually the MSSM parameters:

- M_1, M_2, M_3 : U(1), U(2), U(3) gaugino mass parameters

¹http://www.ihep.ac.cn/lunwen/wjx/public_html

SPS	mSUGRA scenario	m_0	$m_{1/2}$	A_0	$\tan \beta$	sign μ
1	typical point	100	250	-100	10	+
2	focus point region	1450	300	0	10	+
3	model line into coannihilation region	90	400	0	10	+
4	large $\tan \beta$	400	300	0	50	+
5	light stop	150	300	-1000	5	+
6	non-unified gaugino masses $M_1 = 480, M_2 = M_3 = 300$	150	300	0	10	+
	GMSB scenario	Λ	M_{mess}	N_{mess}	$\tan \beta$	sign μ
7	NLSP = $\tilde{\tau}_1$	40,000	80,000	3	15	+
8	NLSP = $\tilde{\chi}_1^0$	100,000	200,000	1	15	+
	AMSB scenario	m_0	$m_{3/2}$		$\tan \beta$	sign μ
9	small $\Delta m(\chi_1^+ - \tilde{\chi}_1^0)$	400	60,000		10	+

Table 1: High-energy scale scenario points corresponding to Snowmass points 1-9[11]

- μ : Higgsino mass parameter
- $\tan \beta$: the ratio of the vev 's of the two-Higgs doublet fields
- A_τ, A_b, A_t : the 3rd generation trilinear couplings
- m_{A^0} : the neutral pseudoscalar Higgs (A^0) mass
- $\tilde{m}_{\tilde{e}_L}^2 (= \tilde{m}_{\tilde{\nu}_{eL}}^2), \tilde{m}_{\tilde{e}_R}^2, \tilde{m}_{\tilde{\mu}_L}^2 (= \tilde{m}_{\tilde{\nu}_{\mu L}}^2), \tilde{m}_{\tilde{\mu}_R}^2, \tilde{m}_{\tilde{u}_L}^2 (= \tilde{m}_{\tilde{d}_L}^2), \tilde{m}_{\tilde{u}_R}^2, \tilde{m}_{\tilde{d}_R}^2, \tilde{m}_{\tilde{c}_L}^2 (= \tilde{m}_{\tilde{s}_L}^2), \tilde{m}_{\tilde{c}_R}^2, \tilde{m}_{\tilde{s}_R}^2$: the SUSY-breaking mass parameters for the 1st/2nd generation sfermions
- $\tilde{m}_{\tilde{\tau}_1}^2, \tilde{m}_{\tilde{\tau}_2}^2, \tilde{m}_{\tilde{b}_2}^2, \tilde{m}_{\tilde{t}_1}^2 (= \tilde{m}_{\tilde{b}_1}^2), \tilde{m}_{\tilde{t}_2}^2$: the SUSY-breaking mass parameters for the 3rd generation sfermions

The Yukawa couplings for the 1st and 2nd generation fermions are fixed to zero as an approximation and the corresponding soft SUSY breaking terms are also fixed to zero in the MSSM model used here. If needed, they can be easily included in our package by FDC.

2 Installation and Usage

2.1 Installation

The package is distributed in a gzip compressed file named *fsdecay.tar.gz* and can be downloaded from FDC homepage:

http://www.ihep.ac.cn/lunwen/wjx/public_html

The CERN program library (CERNLIB) 2003 [12] (or up) and ISAJET are required only for high-energy scale mode. The installation steps are given below and for convenience it is assumed that²:

1. CERNLIB is located at directory */cern/2003*,
2. the current user account is *linuxuser*;
3. the downloaded file *fsdecay.tar.gz* are stored at directory *~linuxuser*;
4. the package will be installed to *~linuxuser/FDCDecay*.

The installation steps are

1. Create the target directory and go into it by

```
mkdir ~linuxuser/FDCDecay
cd ~linuxuser/FDCDecay
```

2. Decompress the downloaded file by

```
tar -xzf ~linuxuser/fsdecay.tar.gz
```

3. Specify the CERNLIB path by adding following line to the **very beginning** of the file *makefile*³

```
cern = /cern/2003
```

4. Compile the low-energy scale mode by

```
make lowscale
```

5. Compile the high-energy scale mode by

```
make highscale
```

6. Delete the temporary files by

```
make clean
```

After steps above without encountering any errors, one will obtain two executable files named *fsdecayhigh* and *fsdecaylow* in directory *~linuxuser/FDCDecay* which are stand-alone programs used to calculate decay width and branch ratio in high-energy scale and low-energy scale modes respectively.

²Users can substitute the user account and paths for their owns, when installing on their own computers.

³This step is necessary only for high-energy scale mode.

2.2 Usage

As mentioned above, with this package, users can calculate decays in high-energy scale mode or low-energy scale mode. While in high-energy scale mode, users can either specify the scenarios (including mSUGRA, GMSB, AMSB) and their parameters manually or just use the Snowmass benchmark points. While in low-energy scale mode, users must input the MSSM (and the Standard Model) parameters manually.

For this purpose, we prepared two executable programs *fsdecayhigh* and *fsdecaylow* corresponding to these two modes respectively.

Next we will describe the usage of these two programs. Test-run examples are given in Sec. 6.

2.2.1 Using high-energy scale mode

It is very simple to calculate decay width in high-energy scale mode. What users need to do is just run program *fsdecayhigh* and then the program will give prompts on screen and require corresponding inputs.

2.2.2 Using low-energy scale mode

To calculate decay width in this mode is somewhat more complicated. Users should follow steps below:

1. Modify the FORTRAN file *usrpara.f* in the installation directory to assign value to parameters. There is a subroutine *FUPAR* in which users can specify the MSSM and SM parameters respectively by assigning values to corresponding variables. All these variables are listed in Tab.2.
2. Re-build the executable program *fsdecaylow* by command:
make lowscale clean
3. Run program *fsdecaylow* and it will give prompt for users to specify the output file name

3 Output files

The format of the generated output file for decays complies with the SUSY Les Houches Accord[9]. It is composed of several blocks which are separated by empty lines. Each of these blocks describes a SUSY particle decay. Every block begins with a DECAF statement specifying the decaying mother particle and its total width, in the format as follows

```
#      PDG      WIDTH(GeV)
DECAF  2000006   9.00247665E+00  # ~t_2 decays
```

MSSM parameters				SM parameters	
FSSGM1	M_1	FSSBMUR	$\tilde{m}_{\tilde{\mu}_R}^2$	FSMME	m_e
FSSGM2	M_2	FSSBTAU1	$\tilde{m}_{\tilde{\tau}_1}^2$	FSMMU	m_μ
FSSGM3	M_3	FSSBTAU2	$\tilde{m}_{\tilde{\tau}_2}^2$	FSMMTAU	m_τ
FSSHMU	μ	FSSBUPL	$\tilde{m}_{\tilde{u}_L}^2$	FSMMUP	m_u
FSS TANB	$\tan \beta$	FSSBUPR	$\tilde{m}_{\tilde{u}_R}^2$	FSMMDN	m_d
FSSMHA	m_{A^0}	FSSBCHL	$\tilde{m}_{\tilde{c}_L}^2$	FSMMCH	m_c
FSSAAL	A_τ	FSSBCHR	$\tilde{m}_{\tilde{c}_R}^2$	FSMMST	m_s
FSSAAT	A_t	FSSBDNR	$\tilde{m}_{\tilde{d}_R}^2$	FSMMTP	m_t
FSSAAB	A_b	FSSBSTR	$\tilde{m}_{\tilde{s}_R}^2$	FSMMBT	m_b
FSSSBEL	$\tilde{m}_{\tilde{e}_L}^2$	FSSBBT2	$\tilde{m}_{\tilde{b}_2}^2$	FSMMZ	m_z
FSSSBER	$\tilde{m}_{\tilde{e}_R}^2$	FSSSBTP1	$\tilde{m}_{\tilde{t}_1}^2$	FSMSN2THW	$\sin^2(\theta_W)$
FSSSBMURL	$\tilde{m}_{\tilde{\mu}_L}^2$	FSSSBTP2	$\tilde{m}_{\tilde{t}_2}^2$	FSMFNST	α_{EM}
				FSMALPS	α_s

Table 2: Variables corresponding to the MSSM and SM parameters in subroutine *FUPAR*

The first integer is the PDG particle code[13] of the decaying particle. The subsequent real number is its total width. The end comment gives a human readable translation of the PDG code. All the decay modes for this mother particle are listed in the subsequent lines in the format

```
#   BR              NDA   ID1      ID2
2.25878090E-01    2      23    1000006 # ~t_2 -> Z0 ~t_1
2.21288383E-01    2          5    1000037 # ~t_2 -> b ~chi+_2
1.90569923E-01    2          6    1000035 # ~t_2 -> t ~chi0_4
1.61424758E-01    2          5    1000024 # ~t_2 -> b ~chi+_1
...
```

In each line above, the first real number is the branch ratio of the decay mode, the first integer is the number of daughters (which is always 2 in this package) and the following integers are the PDG codes of these daughters. The specific FORTRAN format for the DECAY statement and the entries in a block are respectively

```
FORMAT('#',7X,'PDG',9X,'WIDTH(GeV)')
FORMAT('DECAY',1X,I9,3X,1P,E16.8,OP,3X,'#',1X,A25)
FORMAT('#',4X,'BR',16X,'NDA',3X,'ID1',7X,'ID2')
FORMAT(3X,1P,E16.8,OP,3X,I2,3X,2(I9,1X),2X,'#',1X,A25)
```

4 Cross-Check with ISAJET

4.1 Running mass parameters and higher-order mass corrections

In ISAJET (Version 7.64), running mass for τ , b and t are used in the calculations of stau($\tilde{\tau}_i$), sbottom ($\tilde{b}_i$) and stop ($\tilde{t}_i$) mass eigenvalues, mixing angles and in the vertices of $fermion(\tau/b/t) \sim sfermion(\tilde{\tau}_i/\tilde{b}_i/\tilde{t}_i) \sim gaugino(\tilde{\chi}_i^\pm/\tilde{\chi}_i^0)$.

SPS	$M_{H^0}^I$	$M_{H^0}^F$	$M_{h^0}^I$	$M_{h^0}^F$	$M_{H^+}^I$	$M_{H^+}^F$	$M_{\tilde{g}}^I$	$M_{\tilde{g}}^F$
1	395.629	395.310	113.657	89.266	403.229	402.885	606.633	579.093
2	1501.688	1499.753	115.551	89.358	1503.698	1501.773	794.334	691.835
3	581.559	580.898	116.663	89.320	586.758	586.088	932.097	893.408
4	356.491	356.153	114.965	91.092	369.506	364.995	732.394	686.647
5	690.826	691.094	119.901	84.047	695.257	694.805	719.234	687.721
6	464.004	463.524	114.433	89.294	470.489	470.008	717.950	684.761
7	390.072	389.560	113.537	90.317	398.160	397.483	943.832	894.279
8	532.413	531.719	114.802	90.339	538.381	537.554	835.300	753.465
9	926.592	925.408	115.089	89.347	929.825	928.678	1294.129	1225.333

Table 3: Higgs and gluino mass in ISAJET (I) and FDCSUSYDecay (F)

Higher order corrections are added in ISAJET to the mass of Higgs H^0 , h^0 , H^+ and gluino $\tilde{g}$. The corrections at SPS 1-9 bring less than 1% modification to the H^0 and H^+ mass, 27% $\sim$ 43% to h^0 mass and 5% $\sim$ 15% for $\tilde{g}$ mass. In Tab.3 we listed the Higgs and gluino mass at SPS 1-9 from ISAJET and our decay package. However these corrections have not been implemented in FDC yet (thus in our package).

In order to compare the decay widths, we take, **temporarily**, the running mass and the corrected Higgs and gluino mass as inputs from ISAJET.

4.2 Cross-check and results

Taking higher order corrections to Higgs and gluino mass as inputs, we calculated the decay widths at SPS 1-9 and compared them with the data from ISAJET.

For the decay modes with 10% discrepancies or larger between ISAJET and our package, we analyzed the decay equations in the source codes of ISAJET (Version 7.64). Two reasons for the discrepancies were found: (1) correction to triple Higgs vertices and QCD radiative correction to $Higgs \rightarrow q(t/b) + \bar{q}(\bar{t}/\bar{b})$ included in ISAJET; (2) different decay equations used by ISAJET and our package which imply something wrong in the codes of ISAJET⁴.

For the first kind of discrepancy, we list the widths of Higgs decays which are influenced by the higher order corrections in Tab.4. From Tab.4 we can see that the corrections bring large modifications sometimes.

In the following we present a detailed discussion for the second kind of discrepancy. There are 3 kinds of decay modes with different decay equations between ISAJET and FDCSUSYDecay. They are:

1. A chargino decays into a left-sstrange and a c-quark ($\tilde{\chi}_i^+ \rightarrow \bar{s}_L + c$);
2. The neutral pseudoscalar Higgs decays into 2 different staus ($A^0 \rightarrow \tilde{\tau}_1^- + \tilde{\tau}_2^-$);

⁴We carefully checked the MSSM model used in FDC with that in Ref. [14]

$H^0 \rightarrow b + \bar{b}$			$H^0 \rightarrow t + \bar{t}$			$H^0 \rightarrow h^0 + h^0$		
SPS	Γ_I	Γ_F	SPS	Γ_I	Γ_F	SPS	Γ_I	Γ_F
1	1.131	1.956	1	0.0435	0.0300	1	0.00120	0.00945
2	3.621	7.446	2	0.779	0.851	2	0.00275	0.00258
3	1.580	2.879	3	0.226	0.198	3	0.00167	0.00658
4	25.910	44.143	4	1.300×10^{-4}	7.718×10^{-05}	4	0.00155	0.00044
5	0.457	0.854	5	1.262	1.145	5	0.0366	0.01836
6	1.298	2.297	6	0.118	0.0927	6	0.00176	0.00819
7	2.516	4.342	7	0.0175	0.0113	7	0.00416	0.00443
8	3.294	5.933	8	0.0834	0.0691	8	0.00343	0.00332
9	2.371	4.592	9	0.466	0.464	9	0.00275	0.00417
$h^0 \rightarrow b + \bar{b}$			$H^+ \rightarrow t + \bar{b}$					
SPS	Γ_I	Γ_F	SPS	Γ_I	Γ_F			
1	0.00482	0.00685	1	0.835	1.463			
2	0.00405	0.00575	2	3.981	8.1306			
3	0.00445	0.00629	3	1.473	2.696			
4	0.00413	0.00733	4	14.635	26.815			
5	0.00449	0.00626	5	1.697	2.229			
6	0.00462	0.00652	6	1.0817	1.930			
7	0.00509	0.00691	7	1.577	2.938			
8	0.00456	0.00632	8	2.531	4.898			
9	0.00414	0.00586	9	2.484	4.801			

Table 4: Higgs decay widths in ISAJET (I) and FDCSUSYDecay (F)

$A^0 \rightarrow \tilde{\tau}_1^- + \tilde{\tau}_2^+$			$A^0 \rightarrow \chi_1^- + \chi_1^+$		
SPS	Γ_I	Γ_F	SPS	Γ_I	Γ_F
1	0.00442	0.00594	1	0.0519	0.258
2	-	-	2	1.494	1.642
3	0.00351	0.00377	3	-	-
4	-	-	4	-	-
5	0.0254	0.0305	5	0.127	0.223
6	0.00111	0.00132	6	0.0294	0.223
7	2.76×10^{-5}	2.71×10^{-5}	7	-	-
8	3.85×10^{-6}	3.34×10^{-6}	8	0.0177	0.213
9	0.0367	0.130	9	0.0992	0.116
$\chi_2^+ \rightarrow \tilde{s}_L + c$			$A^0 \rightarrow \chi_2^- + \chi_2^+$		
SPS	Γ_I	Γ_F	SPS	Γ_I	Γ_F
5	1.86×10^{-5}	1.7×10^{-4}	2	0.354	0.550

Table 5: Decay width from ISAJET (I) and FDCSUSYDecay (F)

3. The neutral pseudoscalar Higgs decays into a chargino pair ($A^0 \rightarrow \tilde{\chi}_i^- + \tilde{\chi}_i^+$, $i = 1, 2$).

The numerical results of decay widths at SPS 1 to 9 are shown in Tab.5 and it is very clear that there are large discrepancies in these decay channels.

In the following, we briefly present the definition of the particles, the resulting mixing matrices and the related Feynman rules. The mass matrix for stau in the $\tilde{\tau}_L - \tilde{\tau}_R$ basis is

$$M_{\tilde{\tau}} = \begin{pmatrix} \tilde{M}_{\tilde{\tau}_L}^2 & m_{\tau}(A_{\tau} - \mu \tan \beta) \\ m_{\tau}(A_{\tau} - \mu \tan \beta) & \tilde{M}_{\tilde{\tau}_R}^2 \end{pmatrix}$$

$$\tilde{M}_{\tilde{\tau}_L}^2 = m_{\tau}^2 + \tilde{m}_{\tilde{\tau}_1}^2 + m_z^2 \cos 2\beta \left(-\frac{1}{2} + \sin^2 \theta_W\right), \quad (1)$$

$$\tilde{M}_{\tilde{\tau}_R}^2 = m_{\tau}^2 + \tilde{m}_{\tilde{\tau}_2}^2 - m_z^2 \cos 2\beta \sin^2 \theta_W, \quad (2)$$

where $\tilde{m}_{\tilde{\tau}_1}^2$, $\tilde{m}_{\tilde{\tau}_2}^2$ and A_{τ} are parameters in soft SUSY breaking term. The stau mass matrix is diagonalized by a unitary matrix $U^{\tilde{\tau}}$

$$(U^{\tilde{\tau}})^{-1} M_{\tilde{\tau}} U^{\tilde{\tau}} = \begin{pmatrix} m_{\tilde{\tau}_1}^2 & 0 \\ 0 & m_{\tilde{\tau}_2}^2 \end{pmatrix}, \quad (m_{\tilde{\tau}_2} > m_{\tilde{\tau}_1}), \quad (3)$$

and

$$\begin{pmatrix} \tilde{\tau}_L^- \\ \tilde{\tau}_R^- \end{pmatrix} = U^{\tilde{\tau}} \begin{pmatrix} \tilde{\tau}_1^- \\ \tilde{\tau}_2^- \end{pmatrix}, \quad U^{\tilde{\tau}} = \begin{pmatrix} \cos \theta_{\tau} & \sin \theta_{\tau} \\ -\sin \theta_{\tau} & \cos \theta_{\tau} \end{pmatrix}. \quad (4)$$

We adopt the positively charged charginos as particles and define the physical states with mass $m_{\tilde{\chi}_i}$ as

$$\Psi(\tilde{\chi}_i^+) = \begin{pmatrix} \overline{\lambda_{iR}^-} \\ \lambda_{iL}^+ \end{pmatrix}, \quad \Psi(\tilde{\chi}_i^-) \equiv \Psi(\tilde{\chi}_i^+)^C = \begin{pmatrix} \overline{\lambda_{iL}^+} \\ \lambda_{iR}^- \end{pmatrix} \quad (5)$$

where

$$\begin{pmatrix} \lambda^- \\ \tilde{H}_1^- \end{pmatrix} = U \begin{pmatrix} \lambda_{1R}^- \\ \lambda_{2R}^- \end{pmatrix}, \quad \begin{pmatrix} \lambda^+ \\ \tilde{H}_2^+ \end{pmatrix} = V \begin{pmatrix} \lambda_{1L}^+ \\ \lambda_{2L}^+ \end{pmatrix}.$$

The $\lambda^\pm$ denote winos and $\tilde{H}_1^-$ and $\tilde{H}_2^+$ denote charged higgsinos. The unitary matrices U and V are given as

$$U = \begin{pmatrix} \cos \theta_R & \sin \theta_R \\ -\sin \theta_R & \cos \theta_R \end{pmatrix} \begin{pmatrix} \eta_1 & 0 \\ 0 & \eta_2 \end{pmatrix}, \quad V = \begin{pmatrix} \cos \theta_L & \sin \theta_L \\ -\sin \theta_L & \cos \theta_L \end{pmatrix}, \quad (6)$$

The chargino mass matrix is

$$M_{\tilde{\chi}} = \begin{pmatrix} M_2 & \sqrt{2}m_z \cos \theta_W \sin \beta \\ \sqrt{2}m_z \cos \theta_W \cos \beta & \mu \end{pmatrix}. \quad (7)$$

It can be diagonalized by unitary matrices in (6)

$$U^{-1} M_{\tilde{\chi}} V = \begin{pmatrix} \eta_1 m_{\tilde{c}_1} & 0 \\ 0 & \eta_2 m_{\tilde{c}_2} \end{pmatrix}, \quad (|m_{\tilde{c}_2}| > |m_{\tilde{c}_1}|), \quad (8)$$

where $\eta_1 = \text{sign}(m_{\tilde{c}_1})$ and $\eta_2 = \text{sign}(m_{\tilde{c}_2})$. The chargino mass $m_{\tilde{\chi}_i}$ then will be

$$m_{\tilde{\chi}_1} = \eta_1 m_{\tilde{c}_1}, \quad m_{\tilde{\chi}_2} = \eta_2 m_{\tilde{c}_2}. \quad (9)$$

The related Feynman rules used in our package are given in Fig.1 and 2, which have been carefully checked with those in Ref. [14].




Figure 1: Vertices of $\tilde{\chi}_i^+ \bar{s}_L c$ in FDC

$$\begin{aligned}
& \text{Diagram 1: } A_0 \rightarrow \tilde{\tau}_1^- \tilde{\tau}_2^+ \quad \frac{gm_\tau}{2m_W}(\mu + A_\tau \tan \beta) \\
& \text{Diagram 2: } A_0 \rightarrow \tilde{\chi}_1^+ \tilde{\chi}_1^- \quad -\frac{1}{\sqrt{2}}\eta_1 g(\cos \beta \cos \theta_R \sin \theta_L + \sin \beta \sin \theta_R \cos \theta_L) \gamma_5 \\
& \text{Diagram 3: } A_0 \rightarrow \tilde{\chi}_2^+ \tilde{\chi}_2^- \quad \frac{1}{\sqrt{2}}\eta_2 g(\cos \beta \sin \theta_R \cos \theta_L + \sin \beta \cos \theta_R \sin \theta_L) \gamma_5
\end{aligned}$$

Figure 2: Vertices of $A_0 \tilde{\tau}_1^- \tilde{\tau}_2^-$ and $A_0 \tilde{\chi}_i^- \tilde{\chi}_i^+$

and decay width equations are listed as:

$$\Gamma(\tilde{\chi}_1^+ \rightarrow \bar{s}_L + c) = \frac{3g^2|P_{cm}|}{16\pi m_{\tilde{\chi}_1}^2}(m_c^2 - m_{\bar{s}_L}^2 + m_{\tilde{\chi}_1}^2) \cos^2 \theta_R \quad (10)$$

$$\Gamma(\tilde{\chi}_2^+ \rightarrow \bar{s}_L + c) = \frac{3g^2|P_{cm}|}{16\pi m_{\tilde{\chi}_2}^2}(m_c^2 - m_{\bar{s}_L}^2 + m_{\tilde{\chi}_2}^2) \sin^2 \theta_R \quad (11)$$

$$\Gamma(A^0 \rightarrow \tilde{\tau}_1^- + \tilde{\tau}_2^+) = \frac{g^2 m_\tau^2 |P_{cm}|}{32\pi m_{A^0}^2 m_W^2}(\mu + A_\tau \tan \beta)^2 \quad (12)$$

$$\Gamma(A^0 \rightarrow \tilde{\chi}_1^- + \tilde{\chi}_1^+) = \frac{g^2 |P_{cm}|}{8\pi}(\cos \beta \cos \theta_R \sin \theta_L + \sin \beta \sin \theta_R \cos \theta_L)^2 \quad (13)$$

$$\Gamma(A^0 \rightarrow \tilde{\chi}_2^- + \tilde{\chi}_2^+) = \frac{g^2 |P_{cm}|}{8\pi}(\cos \beta \sin \theta_R \cos \theta_L + \sin \beta \cos \theta_R \sin \theta_L)^2 \quad (14)$$

where $|P_{cm}|$ is the momentum of one of the final particle in the rest frame of the initial particle

$$|P_{cm}| = \frac{\sqrt{[M^2 - (m_1 + m_2)^2][M^2 - (m_1 - m_2)^2]}}{2M}$$

where M is the mass of the initial particle and m_1, m_2 is the masses of the final particles.

5 Summary

We have presented the FORTRAN program package FDCSUSYDecay, which is used to calculate all the possible SUSY particle and Higgs 2-body decays in the MSSM model at tree-level. The package with needed 3-body channels can be generated by FDC when the MSSM parameters are fixed. One of the attractive features of this package is that all the FORTRAN codes are generated by FDC with full automation and it can avoid uncaredful errors. Although this package can now only work at tree-level, the ability of one-loop calculation in FDC are under developing. A new version of FDCSUSYDecay with full one-loop correction will be generated in near future.

6 Test Run and Output

6.1 Test-run in high-energy scale mode with parameters of Snowmass benchmark point 1

The screen output when running *fsdecayhigh* is as following:

```
*****
* FDCSUSYDecay 1.00                      *
*****
Enter Snowmass point (1-9):
OR Enter 0 to specify scenario and parameters manually:
1
Enter output file name (in single quotes):
'testrun1.dat'
```

The decay table is stored in file *testrun1.dat* and reads like (partly listed):

```
# =====
#          DECAY TABLE CREATED BY FDCSUSYDECAY
#          DATE: 11/ 4/2006      16:43:46
#          =====
#
#          PDG          WIDTH(GeV)
DECAY  2000006      9.00247665E+00  # ~t_2 decays
#  BR          NDA  ID1  ID2
2.25878090E-01  2      23  1000006  # ~t_2 -> Z0 ~t_1
2.21288383E-01  2      5   1000037  # ~t_2 -> b ~chi+_2
1.90569923E-01  2      6   1000035  # ~t_2 -> t ~chi0_4
1.61424758E-01  2      5   1000024  # ~t_2 -> b ~chi+_1
6.33183903E-02  2      6   1000023  # ~t_2 -> t ~chi0_2
3.98983142E-02  2      6   1000025  # ~t_2 -> t ~chi0_3
3.91332313E-02  2      24  1000005  # ~t_2 -> W+ ~b_1
3.31751880E-02  2      25  1000006  # ~t_2 -> h0 ~t_1
2.53137219E-02  2      6   1000022  # ~t_2 -> t ~chi0_1

#          PDG          WIDTH(GeV)
DECAY  1000021      6.33793298E+00  # ~g decays
#  BR          NDA  ID1  ID2
```

```

1.19224123E-01 2 5 -1000005 # ~g -> b ~b_1bar
1.19224123E-01 2 -5 1000005 # ~g -> bar ~b_1
5.10661138E-02 2 2 -2000002 # ~g -> u ~u_Rbar
5.10661138E-02 2 -2 2000002 # ~g -> ubar ~u_R
5.10542957E-02 2 4 -2000004 # ~g -> c ~c_Rbar
5.10542957E-02 2 -4 2000004 # ~g -> cbar ~c_R
5.08960759E-02 2 1 -2000001 # ~g -> d ~d_Rbar
5.08960759E-02 2 -1 2000001 # ~g -> dbar ~d_R
5.08958197E-02 2 3 -2000003 # ~g -> s ~s_Rbar
5.08958197E-02 2 -3 2000003 # ~g -> sbar ~s_R
3.90571495E-02 2 6 -1000006 # ~g -> t ~t_1bar
3.90571495E-02 2 -6 1000006 # ~g -> tbar ~t_1
3.89890226E-02 2 5 -2000005 # ~g -> b ~b_2bar
3.89890226E-02 2 -5 2000005 # ~g -> bar ~b_2
2.81103874E-02 2 2 -1000002 # ~g -> u ~u_Lbar
2.81103874E-02 2 -2 1000002 # ~g -> ubar ~u_L
2.80978313E-02 2 4 -1000004 # ~g -> c ~c_Lbar
2.80978313E-02 2 -4 1000004 # ~g -> cbar ~c_L
2.13047293E-02 2 1 -1000001 # ~g -> d ~d_Lbar
2.13047293E-02 2 -1 1000001 # ~g -> dbar ~d_L
2.13044513E-02 2 3 -1000003 # ~g -> s ~s_Lbar
2.13044513E-02 2 -3 1000003 # ~g -> sbar ~s_L

```

```

# PDG WIDTH(GeV)
DECAY 1000003 5.00062352E+00 # ~s_L decays
# BR NDA ID1 ID2
6.10919249E-01 2 4 -1000024 # ~s_L -> c ~chi-_1
3.10762902E-01 2 3 1000023 # ~s_L -> s ~chi0_2
3.89466555E-02 2 4 -1000037 # ~s_L -> c ~chi-_2
2.41598639E-02 2 3 1000022 # ~s_L -> s ~chi0_1
1.36997584E-02 2 3 1000035 # ~s_L -> s ~chi0_4
1.51157142E-03 2 3 1000025 # ~s_L -> s ~chi0_3

```

...

6.2 Test-run in high-energy scale mode with manually inputed parameters

The screen output when running *fsdecayhigh* is as following:

```

*****
* FDGSUSYDecay 1.00 *
*****
Enter Snowmass point (1-9):
OR Enter 0 to specify scenario and parameters manually:
0
1.mSUGRA
2.minimal Gauge-Mediated SUSY Breaking (mGMSB)
3.Anomaly-Mediated SUSY Breaking (AMSB)
Please choose scenario (1-3):
1
Enter M_0, M_(1/2), A_0, tan(BETA), sign(MU), M_t(top mass)
100.0 260.0 -100.0 12.0 1.0 175.0
Enter output file name (in single quotes):
'testrun2.dat'

```

The decay table is stored in file *testrun2.dat* and reads like (partly listed):

```
#
# =====
#          DECAY TABLE CREATED BY FDGSUSYDECAY
#          DATE: 11/ 4/2006      16:50:38
#          =====
#
#          PDG          WIDTH(GeV)
DECAY  1000021      6.87489451E+00  # ~g decays
#          BR          NDA   ID1      ID2
1.22817522E-01      2          5   -1000005  # ~g -> b   ~b_1bar
1.22817522E-01      2          -5   1000005  # ~g -> bar ~b_1
5.00789610E-02      2          1   -2000001  # ~g -> d   ~d_Rbar
5.00789610E-02      2          -1   2000001  # ~g -> dbar ~d_R
5.00787337E-02      2          3   -2000003  # ~g -> s   ~s_Rbar
5.00787337E-02      2          -3   2000003  # ~g -> sbar ~s_R
4.98299423E-02      2          2   -2000002  # ~g -> u   ~u_Rbar
4.98299423E-02      2          -2   2000002  # ~g -> ubar ~u_R
4.98194437E-02      2          4   -2000004  # ~g -> c   ~c_Rbar
4.98194437E-02      2          -4   2000004  # ~g -> cbar ~c_R
4.34327716E-02      2          6   -1000006  # ~g -> t   ~t_1bar
4.34327716E-02      2          -6   1000006  # ~g -> tbar ~t_1
3.66751775E-02      2          5   -2000005  # ~g -> b   ~b_2bar
3.66751775E-02      2          -5   2000005  # ~g -> bar ~b_2
2.74132411E-02      2          2   -1000002  # ~g -> u   ~u_Lbar
2.74132411E-02      2          -2   1000002  # ~g -> ubar ~u_L
2.74020784E-02      2          4   -1000004  # ~g -> c   ~c_Lbar
2.74020784E-02      2          -4   1000004  # ~g -> cbar ~c_L
2.12261878E-02      2          1   -1000001  # ~g -> d   ~d_Lbar
2.12261878E-02      2          -1   1000001  # ~g -> dbar ~d_L
2.12259409E-02      2          3   -1000003  # ~g -> s   ~s_Lbar
2.12259409E-02      2          -3   1000003  # ~g -> sbar ~s_L

#          PDG          WIDTH(GeV)
DECAY  2000006      9.48857394E+00  # ~t_2 decays
#          BR          NDA   ID1      ID2
2.19460985E-01      2          5   1000037  # ~t_2 -> b   ~chi+_2
2.12621918E-01      2          23  1000006  # ~t_2 -> Z0  ~t_1
1.97473598E-01      2          6   1000035  # ~t_2 -> t   ~chi0_4
1.65481446E-01      2          5   1000024  # ~t_2 -> b   ~chi+_1
6.29074190E-02      2          6   1000023  # ~t_2 -> t   ~chi0_2
4.22399775E-02      2          6   1000025  # ~t_2 -> t   ~chi0_3
4.13479750E-02      2          24  1000005  # ~t_2 -> W+  ~b_1
3.30474133E-02      2          25  1000006  # ~t_2 -> h0  ~t_1
2.54192679E-02      2          6   1000022  # ~t_2 -> t   ~chi0_1

#          PDG          WIDTH(GeV)
DECAY  1000003      5.17658881E+00  # ~s_L decays
#          BR          NDA   ID1      ID2
6.11686309E-01      2          4   -1000024  # ~s_L -> c   ~chi-_1
3.11939618E-01      2          3   1000023  # ~s_L -> s   ~chi0_2
3.86054484E-02      2          4   -1000037  # ~s_L -> c   ~chi-_2
2.27239525E-02      2          3   1000022  # ~s_L -> s   ~chi0_1
1.35532277E-02      2          3   1000035  # ~s_L -> s   ~chi0_4
1.49144465E-03      2          3   1000025  # ~s_L -> s   ~chi0_3
```

...

6.3 Test-run in low-energy scale mode

The file *usrpara.f* is prepared as:

```
SUBROUTINE FUPAR()
c-----
c      By this subroutine, users can specify the MSSM and SM
c      parameters at low-energy scale.
c
c      ** IMPORTANT **
c      PROGRAM DO NOT CHECK THE VALIDITY OF PARAMETERS VALUE
c      YOU INPUTED.
c
c      **REMEMBER TO RE-BUILD THE PROGRAM AFTER EVERY MODIFICATION
c      BY COMMAND:
c      make lowscale
c-----
c      The MSSM parameter variables are listed below:
c      The unit is GeV
c
c      FSSGM1   = U(1) gaugino mass parameter
c      FSSGM2   = U(2) gaugino mass parameter
c      FSSGM3   = U(3) gaugino mass parameter
c      FSSHMU   = Higgsino mass parameter
c      FSSTANB  = ratio of Higgs vacuum expectation
c      FSSMHA   = neutral pseudoscalar Higgs(A0) mass
c      FSSAAL   = trilinear coupling for tau lepton
c      FSSAAT   = trilinear coupling for t-quark
c      FSSAAB   = trilinear coupling for b-quark
c      FSSSBEL  = SUSY-breaking mass parameter for left s-electron
c      FSSSBER  = SUSY-breaking mass parameter for right s-electron
c      FSSSBMUL = SUSY-breaking mass parameter for left s-muon
c      FSSSBMUR = SUSY-breaking mass parameter for right s-muon
c      FSSSBTAU1 = SUSY-breaking mass parameter for left s-tau
c      FSSSBTAU2 = SUSY-breaking mass parameter for right s-tau
c      FSSSBUPL = SUSY-breaking mass parameter for left s-up
c      FSSSBUPR = SUSY-breaking mass parameter for right s-up
c      FSSSBCHL = SUSY-breaking mass parameter for left s-charm
c      FSSSBCHR = SUSY-breaking mass parameter for right s-charm
c      FSSSBDNR = SUSY-breaking mass parameter for right s-down
c      FSSSBSTR = SUSY-breaking mass parameter for right s-strange
c      FSSSB BT2 = SUSY-breaking mass parameter for heavy s-bottom
c      FSSSBTP1 = SUSY-breaking mass parameter for light s-top
c      FSSSBTP2 = SUSY-breaking mass parameter for heavy s-top
c-----
c      The SM parameter variables are listed below:
c      The unit is GeV
c
c      FSMME    = electron mass
c      FSMMMU   = muon mass
c      FSMMTAU  = tau mass
c      FSMMUP   = u-quark mass
c      FSMDN    = d-quark mass
c      FSMMCH   = c-quark mass
```

```

c      FSMMST      = s-quark mass
c      FSMMTP      = t-quark mass
c      FSMMBT      = b-quark mass
c      FSMMZ       = Z boson mass
c      FSMSN2THW   = weak-mixing angle SIN^2(THETA_W)
c      FSMFNST     = fine-structure constant ALPHA_EM
c      FSMALPS     = strong-coupling constant ALPHA_S
c-----
      IMPLICIT NONE

cc BLOCK FUSSPAR START
      DOUBLE PRECISION FSSGM1, FSSGM2,FSSGM3,FSSHMU,FSSTANB,FSSMHA,
$      FSSAAL,FSSAAT,FSSAAB,FSSSBEL,FSSSBER,FSSSBMUL,FSSSBMUR,
$      FSSSBTAU1,FSSSBTAU2,FSSSBUPL,FSSSBUPR,FSSSBCHL,FSSSBCHR,
$      FSSBDNR,FSSSBSTR,FSSBBT2,FSSSBTP1,FSSSBTP2

      COMMON /FUSSPAR/ FSSGM1, FSSGM2,FSSGM3,FSSHMU,FSSTANB,FSSMHA,
$      FSSAAL,FSSAAT,FSSAAB,FSSSBEL,FSSSBER,FSSSBMUL,FSSSBMUR,
$      FSSSBTAU1,FSSSBTAU2,FSSSBUPL,FSSSBUPR,FSSSBCHL,FSSSBCHR,
$      FSSBDNR,FSSSBSTR,FSSBBT2,FSSSBTP1,FSSSBTP2
cc BLOCK END

cc BLOCK FUSMPAR START
      DOUBLE PRECISION FSMME,FSMMMU,FSMMTAU,FSMMUP,FSMMDN,FSMMCH,
$      FSMMST,FSMMBT,FSMMTP,FSMMZ,FSMSN2THW,FSMFNST,FSMALPS

      COMMON /FUSMPAR/ FSMME,FSMMMU,FSMMTAU,FSMMUP,FSMMDN,FSMMCH,
$      FSMMST,FSMMBT,FSMMTP,FSMMZ,FSMSN2THW,FSMFNST,FSMALPS
cc BLOCK END

cc SET YOUR MSSM PARAMETERS HERE

      FSSGM1      = 99.6476898d0
      FSSGM2      = 192.514954d0
      FSSGM3      = 579.09314d0
      FSSHMU      = 354.201385d0
      FSSTANB     = 10.0D0
      FSSMHA      = 394.874817D0
      FSSAAL      = -253.15593d0
      FSSAAB      = -766.899109d0
      FSSAAT      = -495.786896d0
      FSSSBEL     = 38581.48830D0
      FSSSBER     = 18547.65040D0
      FSSSBMUL    = 38581.48830D0
      FSSSBMUR    = 18547.65040D0
      FSSSBTAU1   = 38243.67190D0
      FSSSBTAU2   = 17847.84380D0
      FSSSBUPL    = 289769.750D0
      FSSSBUPR    = 271373.5620D0
      FSSSBCHL    = 289769.750D0
      FSSSBCHR    = 271373.5620D0
      FSSBDNR     = 269598.8440D0
      FSSSBSTR    = 269598.8440D0
      FSSBBT2     = 266551.1560D0
      FSSSBTP1    = 243069.0310D0
      FSSSBTP2    = 178026.6410D0

```


cc SET YOUR SM PARAMETERS HERE

```

FSMME      = 5.10999991D-4
FSMMMU     = 1.04999997D-1
FSMMTAU    = 1.77699995
FSMMUP     = 5.5999998D-3
FSMMDN     = 9.89999995D-3
FSMMCH     = 1.35000002D0
FSMMST     = 1.99000001D-1
FSMMTP     = 175D0
FSMMBT     = 5D0
FSMMZ      = 91.1699982D0
FSMSN2THW  = 0.2311678d0
FSMFNST    = 1D0/127.970535
FSMALPS    = 0.118170463d0
END

```

The screen output when running *fsdecaylow* is as following:

```

*****
* FDCSUSYDecay 1.00                      *
*****
ENTER output file name (in single quotes):
'testrun3.dat'

```

The decay table is stored in file *testrun3.dat* and reads like (partly listed):

```

#          =====
#          DECAY TABLE CREATED BY FDCSUSYDECAY
#          DATE: 11/ 4/2006    16:58:46
#          =====
###
#          PDG          WIDTH(GeV)
DECAY  2000006      9.00247595E+00  # ~t_2 decays
#          BR          NDA    ID1    ID2
2.25878090E-01    2          23    1000006  # ~t_2 -> Z0 ~t_1
2.21288384E-01    2          5     1000037  # ~t_2 -> b ~chi+_2
1.90569924E-01    2          6     1000035  # ~t_2 -> t ~chi0_4
1.61424752E-01    2          5     1000024  # ~t_2 -> b ~chi+_1
6.33183886E-02    2          6     1000023  # ~t_2 -> t ~chi0_2
3.98983144E-02    2          6     1000025  # ~t_2 -> t ~chi0_3
3.91332354E-02    2          24    1000005  # ~t_2 -> W+ ~b_1
3.31751871E-02    2          25    1000006  # ~t_2 -> h0 ~t_1
2.53137244E-02    2          6     1000022  # ~t_2 -> t ~chi0_1

#          PDG          WIDTH(GeV)
DECAY  1000021      6.33793301E+00  # ~g decays
#          BR          NDA    ID1    ID2
1.19224123E-01    2          5    -1000005  # ~g -> b ~b_1bar
1.19224123E-01    2          -5    1000005  # ~g -> bar ~b_1
5.10661146E-02    2          2    -2000002  # ~g -> u ~u_Rbar
5.10661146E-02    2          -2    2000002  # ~g -> ubar ~u_R
5.10542966E-02    2          4    -2000004  # ~g -> c ~c_Rbar
5.10542966E-02    2          -4    2000004  # ~g -> cbar ~c_R
5.08960753E-02    2          1    -2000001  # ~g -> d ~d_Rbar
5.08960753E-02    2          -1    2000001  # ~g -> dbar ~d_R
5.08958191E-02    2          3    -2000003  # ~g -> s ~s_Rbar
5.08958191E-02    2          -3    2000003  # ~g -> sbar ~s_R

```

3.90571491E-02	2	6	-1000006	# $\tilde{g} \rightarrow t \tilde{t}_1\text{bar}$
3.90571491E-02	2	-6	1000006	# $\tilde{g} \rightarrow t\text{bar} \tilde{t}_1$
3.89890229E-02	2	5	-2000005	# $\tilde{g} \rightarrow b \tilde{b}_2\text{bar}$
3.89890229E-02	2	-5	2000005	# $\tilde{g} \rightarrow b\text{bar} \tilde{b}_2$
2.81103873E-02	2	2	-1000002	# $\tilde{g} \rightarrow u \tilde{u}_L\text{bar}$
2.81103873E-02	2	-2	1000002	# $\tilde{g} \rightarrow u\text{bar} \tilde{u}_L$
2.80978312E-02	2	4	-1000004	# $\tilde{g} \rightarrow c \tilde{c}_L\text{bar}$
2.80978312E-02	2	-4	1000004	# $\tilde{g} \rightarrow c\text{bar} \tilde{c}_L$
2.13047294E-02	2	1	-1000001	# $\tilde{g} \rightarrow d \tilde{d}_L\text{bar}$
2.13047294E-02	2	-1	1000001	# $\tilde{g} \rightarrow d\text{bar} \tilde{d}_L$
2.13044514E-02	2	3	-1000003	# $\tilde{g} \rightarrow s \tilde{s}_L\text{bar}$
2.13044514E-02	2	-3	1000003	# $\tilde{g} \rightarrow s\text{bar} \tilde{s}_L$

#	PDG	WIDTH(GeV)		# $\tilde{s}_L$ decays	
DECAY	1000003	5.00062300E+00			
#	BR	NDA	ID1	ID2	
	6.10919249E-01	2	4	-1000024	# $\tilde{s}_L \rightarrow c \tilde{\chi}_{-1}$
	3.10762900E-01	2	3	1000023	# $\tilde{s}_L \rightarrow s \tilde{\chi}_{0_2}$
	3.89466552E-02	2	4	-1000037	# $\tilde{s}_L \rightarrow c \tilde{\chi}_{-2}$
	2.41598663E-02	2	3	1000022	# $\tilde{s}_L \rightarrow s \tilde{\chi}_{0_1}$
	1.36997584E-02	2	3	1000035	# $\tilde{s}_L \rightarrow s \tilde{\chi}_{0_4}$
	1.51157143E-03	2	3	1000025	# $\tilde{s}_L \rightarrow s \tilde{\chi}_{0_3}$

...

References

- [1] E. Boose *et al.*, INP MSU-94-36/358, SNUTP 94-116 (Seoul, 1994); hep-ph/9503280
- [2] T. Ishikawa *et al.*, Grace manual, KEK report 92-19(1993), F. Yuasa *et al.*, Prog. Theor. Phys. Suppl. **138**,18 (2000).
- [3] J.X. Wang, Nucl. Instrum. Meth. A **534**, 241 (2004) [arXiv:hep-ph/0407058].
- [4] J. X. Wang, Prepared for 6th Accelerator and Particle Physics Institute (APPI 2001), Morioka, Japan, 20-23 Feb 2001. Published in Appi 2001, “Accelerator and particle physics,” 108-121
- [5] W. Qi and J. X. Wang, In preparing.
- [6] A. Djouadi, *et al.*, Comput. Phys. Commun. **108**(1998) 56.
- [7] M. Muhlleitner, *et al.*, Comput. Phys. Commun. 168(2005) 46.
- [8] M. Battaglia, *et al.*, Eur. Phys. J. **C25**, 113 (2002), [arXiv:hep-ph/0202233].
- [9] P. Skands *et al.*, JHEP **0407**, 036 (2004) [arXiv:hep-ph/0311123].
- [10] H. Baer, *et al.*, “ISAJET 7.69: A Monte Carlo Event Generator for pp, $\bar{p}p$, and e^+e^- Reactions,” arXiv:hep-ph/0312045.
- [11] N. Ghodbane and H. U. Martyn, “Compilation of SUSY particle spectra from Snowmass 2001 benchmark,” arXiv:hep-ph/0201233.
- [12] CERN Program Library, URL:<http://cernlib.web.cern.ch/cernlib/>
- [13] Particle Data Group, Physics Letters **B592**, 1 (2004).
- [14] M. Kuroda, “Complete Lagrangian of MSSM,” arXiv:hep-ph/9902340;
H. E. Haber and G. L. Kane, “The Search For Supersymmetry: Probing Physics Beyond The Standard Model,” Phys. Rept. 117, 75 (1985);
J. Gunion, H. Haber, G. Kane and S. Dawson, “The Higgs Hunter’s Guide, Addison-Wesley,” New York (1990).